Published by :
http://www.ijert.org

International Journal of Engineering Research & Technology (IJERT)
ISSN: 2278-0181
Volume 13, Issue 12, December 2024

Integral Involving H-Function of Several Complex Variables and Exponential Function

Integral Involving H-function and Exponential function

Naresh Bhati
Department of Mathematics and Statistics
Jai Narain Vyas University,
Jodhpur, India

Rajeev Kumar Gupta
Department of Mathematics and Statistics
Jai Narain Vyas University,
Jodhpur, India, Country

***Abstract*— In this paper we have integrated a H-function of several complex variables with combination of hypergeometric function and an exponential function as product. To perform the integrations, we have used the definite integral listed in "Table of Integrals, Series, and Products" by Gradshteyn and Ryzhik. The integrations obtain in this paper will be useful to solve problems of mathematical, statistical and physical sciences where different kind of functions occurs as product. The result obtained from integration was put in compact form.**

***Keywords*— Multivariable H-function, Exponential function, Hypergeometric function.**

1. INTRODUCTION

The German mathematician Carl Friedrich Gauss (1777-1855) introduced a series called gauss hypergeometric series and represent by

$$ {}_2F_1(\alpha,\beta;\gamma;x)=\sum_{u=0}^{\infty}\frac{(\alpha)_u(\beta)_u}{(\gamma)_u}\frac{x^u}{u!} \qquad (1)$$

where is the Pochhammer symbol defined by

$$ (\alpha)_u=\begin{cases}\alpha(\alpha+1)\ldots(\alpha+u-1) & u\in N\\ 1 & u=1\end{cases} \qquad (2)$$

The hypergeometric function is defined for |z| < 1, provided that γ is neither zero nor a negative integer. The gaussian hypergeometric function has fundamental importance in theory of special functions. The most of functions used in physical sciences, applied mathematics, engineering etc., are expressible as its special cases.

The H-function of several complex variables, defined H. M. Srivastava and R. Panda in series of their research papers [5]. The H-function defined and represented in terms of a multiple Mellin-Bernes type contour integral as

$$ H[z_1,z_2,\ldots,z_r]=H^{M,N;m_1,n_1;\ldots;m_r,n_r}_{P,Q;p_1,q_1;\ldots;p_r,q_r}\left[\begin{matrix}z_1\\ \vdots\\ z_r\end{matrix}\middle|\begin{matrix}(a_j;A_j^{(1)},A_j^{(2)},\ldots A_j^{(r)})_{1,P};(c_j^{(1)},C_j^{(1)})_{1,p_1};\\ (b_j,B_j^{(1)},B_j^{(2)},\ldots,B_j^{(r)})_{1,Q};(d_j^{(1)},D_j^{(1)})_{1,q_1};\end{matrix}\right. \qquad (3)$$

$$ \left.\begin{matrix}\ldots;(c_j^{(r)},C_j^{(r)})_{1,p_r}\\ \ldots;(d_j^{(r)},D_j^{(r)})_{1,q_r}\end{matrix}\right]=\frac{1}{(2\pi i)^r}\int_{\Upsilon_1}\ldots\int_{\Upsilon_r}\psi(s_1,\ldots,s_r)\left\{\prod_{k=0}^{r}\phi_k(s_k)z_k^{s_k}ds_k\right\}$$

denote the H-function of r complex variables $z_1,z_2,\ldots,z_r$. Where $i=(-1)^{1/2}$ and

$$ \psi(s_1,s_2,\ldots,s_r)=\frac{\prod_{j=i}^{M}\Gamma\left(b_j-\sum_{k=1}^{r}B_j^{(k)}s_k\right)\prod_{j=1}^{N}\Gamma\left(1-a_j+\sum_{k=1}^{r}A_j^{(k)}s_k\right)}{\prod_{j=M+1}^{Q}\Gamma\left(1-b_j+\sum_{k=1}^{r}B_j^{(k)}s_k\right)\prod_{j=N+1}^{P}\Gamma\left(a_j-\sum_{k=1}^{r}A_j^{(k)}s_k\right)} \qquad (4)$$

$$ \phi_k(s_k)=\frac{\prod_{j=1}^{m_k}\Gamma\left(d_j^{(k)}-D_j^{(k)}s_k\right)\prod_{j=1}^{n_k}\Gamma\left(1-c_j^{(k)}+C_j^{(k)}s_k\right)}{\prod_{j=m_k+1}^{q_k}\Gamma\left(1-d_j^{(k)}+D_j^{(k)}s_k\right)\prod_{j=n_k+1}^{p_k}\Gamma\left(c_j^{(k)}-C_j^{(k)}s_k\right)} \qquad (5)$$

where $(k=1,2,\ldots,r)$

$\begin{cases}a_j,[j=1,2,\ldots,P];c_j^{(k)},[j=1,2,\ldots,p_i;\forall k\in\{1,2,\ldots,r\}]\\ b_j,[j=1,2,\ldots,Q];d_j^{(k)},[j=1,2,\ldots,q_i;\forall i\in\{1,2,\ldots,r\}]\end{cases}$ are complex numbers, and their corresponding related coefficients $\begin{cases}A_j^{(k)},[j=1,2,\ldots,P];C_j^{(k)}[j=1,2,\ldots,p_k;\forall k\in\{1,2,\ldots,r\}]\\ B_j^{(k)},[j=1,2,\ldots,Q];D_j^{(k)}[j=1,2,\ldots,q_k;\forall k\in\{1,2,\ldots,r\}]\end{cases}$ are positive real numbers.

And Υ_k, represent the contours start at the point $\vartheta_k-i\infty$ and goes to the point $\vartheta_k+i\infty$ with $\vartheta_k\in R$, $(k=1,2,\ldots,r)$. The integral in (3) is converges absolutely, under the conditions Srivastava et al. [5] if

$$ |\arg z_k|<\frac{\pi}{2}\tau_k. \qquad (k=1,2,\ldots,r) \qquad (6)$$

Where $\Delta_k\equiv\sum_{j=1}^{P}A_j^{(k)}+\sum_{j=1}^{p_k}C_j^{(k)}-\sum_{j=1}^{Q}B_j^{(k)}-\sum_{j=1}^{q_k}D_j^{(k)}\le 0$

$$ \tau_k=\sum_{j=1}^{N}A_j^{(k)}-\sum_{j=N+1}^{P}A_j^{(k)}+\sum_{j=1}^{M}B_j^{(k)}-\sum_{j=M+1}^{Q}B_j^{(k)}+\sum_{j=1}^{n_k}c_j^{(k)}-\sum_{j=n_k+1}^{p_k}C_j^{(k)}+\sum_{j=1}^{m_k}D_j^{(k)}-\sum_{j=m_k+1}^{q_k}D_j^{(k)}>0 \qquad (7)$$

$,\forall(k=1,2,\ldots,r)$

(This work is licensed under a Creative Commons Attribution 4.0 International License.)

International Journal of Engineering Research & Technology (IJERT)
ISSN: 2278-0181
Volume 13, Issue 12, December 2024

where M,N,P,Q,m_k,n_k,p_k,q_k are positive integer and restricted by the $0\le N\le P$, $Q\ge M\ge 0$, and $q_k\ge m\ge 0$, $p_k\ge n_k\ge 0$, $\forall k\in\{1,2,...,r\}$ and inequalities (6) suitably constrained values of the complex variables $z_1,z_2,...,z_r$. The points $z_k=0, k=1,2,..,r$ and many exceptional parameter values, being tacitly excluded. From Srivastava and Panda [5], we have

$$H[z_1,z_2,...,z_r]=o\left(|z_1|^{e_1}....|z_r|^{e_r}\right)\left(\lim_{1\le j\le m_r}\|z_j\|\to 0\right),\quad \text{where}$$

$$e_k=\lim_{1\le j\le m_r}\operatorname{Re}\left(\frac{d_j^{(k)}}{D_j^{(k)}}\right)\ (k=1,2,...,r) \tag{8}$$

Lemma:
From the table of integration, series and products I.S. Gradshteyn, M.I. Ryzhik [3, (2007): Eq. 3.194 (7), p.316] We need the following integration formula

$$\int_0^\infty x^m(a+bx)^{-n}dx=\frac{a^{m-n+1}}{b^{m+1}}\frac{\Gamma(m+1)\Gamma(n-m-1)}{\Gamma(n)} \tag{9}$$

$[n>m+1>0, a>0; b>0]$

2. MAIN RESULTS:

In this section we have obtain some integrals involving the product of the basic hypergeometric function with H-function of several complex variable and exponential function.

Theorem 1:
If we take $\rho>\sigma+1>0$; $a>0; b>0$; $\xi\ge 0,\delta\ge 0$; $\lambda\ge 0;\eta\ge 0$; *and* $\mu_i\ge o,\upsilon_i\ge 0$, $(\forall i=1,2,...,r)$ *then the following integration hold*

$$\int_0^\infty y^\sigma(a+by)^{-\rho}e^{\delta y^\xi}\,{}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda(a+by)^\eta\right]\times$$
$$H^{M,N;m_1,n_1;...;m_r,n_r}_{P,Q;p_1,q_1;...,p_r,q_r}\left[\begin{matrix}c_1z_1y^{\mu_1}(a+by)^{\upsilon_1}\\ \vdots\\ c_rz_ry^{\mu_r}(a+by)^{\upsilon_r}\end{matrix}\right]dx=\frac{a^{\sigma-\rho+1}}{b^{\sigma+1}}e^{\delta\left(\frac{a}{b}\right)^\xi}\,{}_2F_1\left(\alpha,\beta;\gamma;\frac{ca^{\lambda+\eta}}{b^\lambda}\right)\times$$
$$\sum_{k=o}^\infty\sum_{u=o}^\infty H^{M+1,N+1;m_1,n_1;...;m_r,n_r}_{P+2,Q+1;p_1,q_1;...,p_r,q_r}\left[\begin{matrix}a^{\mu_1+\upsilon_1}b^{-\mu_1}c_1z_1\\ \\ a^{\mu_r+\upsilon_r}b^{-\mu_r}c_rz_r\end{matrix}\left|\begin{matrix}(-\sigma-u\lambda-k\xi;\mu_1,..,\mu_r);\\ \begin{pmatrix}\rho-\eta u-\sigma-u\lambda-k\xi-1;\\ \upsilon_1+\mu_1,..,\upsilon_r+\mu_r\end{pmatrix};\end{matrix}\right.\right. \tag{10}$$
$$\left.\begin{matrix}(a_j;A_j^{(1)},A_j^{(2)},...A_j^{(r)})_{2,P+1};(\rho-u\eta;\upsilon_1,..,\upsilon_r);(c_j^{(1)},C_j^{(1)})_{1,p_1};...;(c_j^{(r)},C_j^{(r)})_{1,p_r}\\ (b_j,B_j^{(1)},B_j^{(2)},...,B_j^{(r)})_{1,Q+1};(d_j^{(1)},D_j^{(1)})_{1,q_1};...;(d_j^{(r)},D_j^{(r)})_{1,q_r}\end{matrix}\right]$$

The above integral will be convergence for condition (6), (7) and (8).

PROOF:

$$\int_0^\infty y^\sigma(a+by)^{-\rho}e^{\delta y^\xi}\,{}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda(a+by)^\eta\right]\times$$
$$H\left[c_1z_1y^{\mu_1}(a+by)^{\upsilon_1},...,c_rz_ry^{\mu_r}(a+by)^{\upsilon_r}\right]dy$$

Now we replace $e^{\delta y^\xi}$ by $\sum_{k=0}^\infty\frac{(\delta y^\xi)^k}{k!}$ and express the hypergeometric function (1) and H-function of several complex variables (3), then we get

$$=\int_0^\infty y^\sigma(a+by)^{-\rho}\sum_{k=0}^\infty\frac{\delta^k y^{k\xi}}{k!}\sum_{u=0}^\infty\frac{(\alpha)_u(\beta)_u c^u y^{u\lambda}(a+by)^{u\eta}}{(\gamma)_u u!(2\pi i)^r}\int_{\Upsilon_1}...\int_{\Upsilon_r}\psi(s_1,...,s_r)\times$$
$$\phi_1(s_1)...\phi_r(s_r)c_1^{S_1}z_1^{S_1}y^{\mu_1S_1}(a+by)^{\upsilon_1S_1}...c_r^{S_r}z_1^{S_1}y^{\mu_1S_1}(a+by)^{\upsilon_1S_1}dS_1..dS_rdy$$

$$=\int_0^\infty y^\sigma(a+by)^{-\rho}\sum_{k=0}^\infty\sum_{u=0}^\infty\frac{\delta^k c^u(\alpha)_u(\beta)_u y^{u\lambda+k\xi}(a+by)^{u\eta}}{(\gamma)_u k!u!(2\pi i)^r}\int_{\Upsilon_1}...\int_{\Upsilon_r}\psi(s_1,...,s_r)\times$$
$$\phi_1(s_1)...\phi_r(s_r)c_1^{S_1}z_1^{S_1}y^{\mu_1S_1}(a+by)^{\upsilon_1S_1}...c_r^{S_r}z_r^{S_r}y^{\mu_rS_r}(a+by)^{\upsilon_rS_r}dS_1...dS_rdy$$

By interchanging the order of integration and summation, we get

$$=\sum_{k=0}^\infty\sum_{u=0}^\infty\frac{c^u\delta^k(\alpha)_u(\beta)_u}{(\gamma)_u k!u!(2\pi i)^r}\int_{\Upsilon_1}...\int_{\Upsilon_r}\psi(s_1,...,s_r)\phi_1(s_1)...\phi_r(s_r)\times$$
$$\left\{\int_0^\infty y^{\sigma+u\lambda+k\xi+\sum_{i=1}^r\mu_iS_i}(a+by)^{\eta u-\rho+\sum_{i=1}^r\upsilon_iS_i}dy\right\}c_1^{S_1}z_1^{S_1}...c_r^{S_r}z_1^{S_1}dS_1..dS_r$$

$$=\sum_{k=0}^\infty\sum_{u=0}^\infty\frac{c^u\delta^k(\alpha)_u(\beta)_u}{(\gamma)_u k!u!(2\pi i)^r}\int_{\Upsilon_1}...\int_{\Upsilon_r}\frac{a^{\sigma+u\lambda+k\xi+\sum_{i=1}^r\mu_iS_i+\eta u-\rho+\sum_{i=1}^r\upsilon_iS_i+1}}{b^{\sigma+u\lambda+k\xi+\sum_{i=1}^r\mu_iS_i+1}}$$
$$\frac{\Gamma\left(\sigma+u\lambda+k\xi+\sum_{i=1}^r\mu_iS_i+1\right)\Gamma\left\{\rho-\eta u-\sum_{i=1}^r\upsilon_iS_i-\sigma-u\lambda-k\xi-\sum_{i=1}^r\mu_iS_i-1\right\}}{\Gamma\left(\rho-\eta u-\sum_{i=1}^r\upsilon_iS_i\right)}$$
$$\psi(s_1,...,s_r)\phi_1(s_1)...\phi_r(s_r)c_1^{S_1}z_1^{S_1}...c_r^{S_r}z_1^{S_1}dS_1..dS_r$$

$$=\frac{a^{\sigma-\rho+1}}{b^{\sigma+1}}\sum_{k=0}^\infty\sum_{u=0}^\infty\frac{c^u\delta^k(\alpha)_u(\beta)_u a^{u\lambda+\eta u+k\xi}}{(\gamma)_u b^{k\xi+u\lambda}k!u!(2\pi i)^r}\times\int_{\Upsilon_1}...\int_{\Upsilon_r}\frac{a^{\sum_{i=1}^r\mu_iS_i+\sum_{i=1}^r\upsilon_iS_i}}{b^{\sum_{i=1}^r\mu_iS_i}}\times$$
$$\frac{\Gamma\left(\sigma+u\lambda+k\xi+\sum_{i=1}^r\mu_iS_i+1\right)\Gamma\left(\rho-\eta u-\sigma-u\lambda-k\xi-\sum_{i=1}^r\mu_iS_i-\sum_{i=1}^r\upsilon_iS_i-1\right)}{\Gamma\left(\rho-\eta u-\sum_{i=1}^r\upsilon_iS_i\right)}$$
$$\psi(s_1,...,s_r)\phi_1(s_1)...\phi_r(s_r)c_1^{S_1}z_1^{S_1}...c_r^{S_r}z_r^{S_r}dS_1..dS_r$$

by virtue, interpreting of equation (1) and (3), we obtain the required result.

$$=\frac{a^{\sigma-\rho+1}}{b^{\sigma+1}}e^{\delta\left(\frac{a}{b}\right)^\xi}\,{}_2F_1\left(\alpha,\beta;\gamma;\frac{ca^{\lambda+\eta}}{b^\lambda}\right)\sum_{k=o}^\infty\sum_{u=o}^\infty H^{M+1,N+1;m_1,n_1;...;m_r,n_r}_{P+2,Q+1;p_1,q_1;...,p_r,q_r}$$
$$\left[\begin{matrix}a^{\mu_1+\upsilon_1}b^{-\mu_1}c_1z_1\\ \\ a^{\mu_r+\upsilon_r}b^{-\mu_r}c_rz_r\end{matrix}\left|\begin{matrix}(-\sigma-u\lambda-k\xi;\mu_1,..,\mu_r);(a_j;A_j^{(1)},A_j^{(2)},...A_j^{(r)})_{2,P+1};\\ (\rho-\eta u-\sigma-u\lambda-k\xi-1;\upsilon_1+\mu_1,..,\upsilon_r+\mu_r);\end{matrix}\right.\right.$$
$$\left.\begin{matrix}(\rho-u\eta;\upsilon_1,..,\upsilon_r);(c_j^{(1)},C_j^{(1)})_{1,p_1};...;(c_j^{(r)},C_j^{(r)})_{1,p_r}\\ (b_j,B_j^{(1)},B_j^{(2)},...,B_j^{(r)})_{1,Q+1};(d_j^{(1)},D_j^{(1)})_{1,q_1};...;(d_j^{(r)},D_j^{(r)})_{1,q_r}\end{matrix}\right]$$

Hence theorem 1 is proved.

Theorem 2:
If we take $\rho>\sigma+1>0$; $a>0;b>0$; $\xi\ge 0,\delta\ge 0$; $\lambda\ge 0;\eta\ge 0$ *and* $\mu_i\ge o,\upsilon_i\ge 0$; $(\forall i=1,2,...,r)$ *then the following integration hold*

$$\int_0^\infty y^\sigma(a+by)^{-\rho}e^{\delta y^\xi}\,{}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda(a+by)^\eta\right]\times$$
$$H^{M,N;m_1,n_1;...;m_r,n_r}_{P,Q;p_1,q_1;...,p_r,q_r}\left[\begin{matrix}c_1z_1y^{-\mu_1}(a+by)^{-\upsilon_1}\\ \vdots\\ c_rz_ry^{-\mu_r}(a+by)^{-\upsilon_r}\end{matrix}\right]dx=\frac{a^{\sigma-\rho+1}}{b^{\sigma+1}}e^{\delta\left(\frac{a}{b}\right)^\xi}\,{}_2F_1\left(\alpha,\beta;\gamma;\frac{ca^{\lambda+\eta}}{b^\lambda}\right)$$
$$\sum_{k=o}^\infty\sum_{u=o}^\infty H^{M+1,N+1;m_1,n_1;...;m_r,n_r}_{P+1,Q+2;p_1,q_1;...,p_r,q_r}\left[\begin{matrix}a^{-\mu_1-\upsilon_1}b^{\mu_1}c_1z_1\\ \\ a^{-\mu_r-\upsilon_r}b^{\mu_r}c_rz_r\end{matrix}\left|\begin{matrix}\begin{pmatrix}2+\sigma+k\xi+u\lambda+u\eta-\rho;\\ \mu_1+\upsilon_1,...,\mu_r+\upsilon_r\end{pmatrix};\\ (1+\sigma+k\xi+u\lambda;\mu_1,..,\mu_r);\end{matrix}\right.\right. \tag{11}$$
$$\left.\begin{matrix}(a_j;A_j^{(1)},A_j^{(2)},...A_j^{(r)})_{2,P+1};(c_j^{(1)},C_j^{(1)})_{1,p_1};...;(c_j^{(r)},C_j^{(r)})_{1,p_r}\\ (b_j,B_j^{(1)},B_j^{(2)},...,B_j^{(r)})_{2,Q+1};(1-\rho+u\eta;\upsilon_1,..,\upsilon_r);(d_j^{(1)},D_j^{(1)})_{1,q_1};...;(d_j^{(r)},D_j^{(r)})_{1,q_r}\end{matrix}\right]$$

The above integral will be convergence for condition (6), (7) and (8).

(This work is licensed under a Creative Commons Attribution 4.0 International License.)

International Journal of Engineering Research & Technology (IJERT)
ISSN: 2278-0181
Volume 13, Issue 12, December 2024

Theorem 3:
If we take $\rho > \sigma + 1 > 0;\quad a > 0; b > 0;\quad \xi \geq 0, \delta \geq 0;\quad \lambda \geq 0; \eta \geq 0$ *and* $\mu_i \geq o, \upsilon_i \geq 0;\quad (\forall i = 1,2,...,r)$ *then the following integration hold*

$$\int_0^\infty y^\sigma (a+by)^{-\rho} e^{\delta y^\xi}\, {}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda (a+by)^\eta\right]\times$$
$$H^{M,N;m_1,n_1;...;m_r,n_r}_{P,Q;p_1,q_1;...;p_r,q_r}\left[\begin{matrix} c_1 z_1 y^{-\mu_1}(a+by)^{\upsilon_1} \\ \vdots \\ c_r z_r y^{-\mu_r}(a+by)^{\upsilon_r}\end{matrix}\right] dx = \frac{a^{\sigma-\rho+1}}{b^{\sigma+1}} e^{\delta\left(\frac{a}{b}\right)^\xi}\, {}_2F_1\left(\alpha,\beta;\gamma;\frac{ca^{\lambda+\eta}}{b^\lambda}\right)$$
$$\sum_{k=o}^\infty\sum_{u=0}^\infty H^{M+1,N+1;m_1,n_1;...;m_r,n_r}_{P+2,Q+1;p_1,q_1;...;p_r,q_r}\left[\begin{matrix} a^{\upsilon_1-\mu_1} b^{\mu_1} c_1 z_1 \\ a^{\upsilon_r-\mu_r} b^{\mu_r} c_r z_r\end{matrix}\middle|\begin{matrix}\begin{pmatrix}2+\sigma+k\xi+u\lambda+u\eta-\rho; \\ \mu_1-\upsilon_1,...,\mu_r-\upsilon_r\end{pmatrix}; \\ (1+\sigma+k\xi+u\lambda;\mu_1,...,\mu_r);\end{matrix}\right. \tag{12}$$
$$\left.\begin{matrix}\left(a_j;A_j^{(1)},A_j^{(2)},...A_j^{(r)}\right)_{2,P+1};\left(\rho-u\eta;\upsilon_1,..,\upsilon_r\right);\left(c_j^{(1)},C_j^{(1)}\right)_{1,p_1};...;\left(c_j^{(r)},C_j^{(r)}\right)_{1,p_r} \\ \left(b_j,B_j^{(1)},B_j^{(2)},...,B_j^{(r)}\right)_{2,Q+1};\left(d_j^{(1)},D_j^{(1)}\right)_{1,q_1};...;\left(d_j^{(r)},D_j^{(r)}\right)_{1,q_r}\end{matrix}\right]$$

The above integral will be convergence for condition (6), (7) and (8).

Theorem 4:
If we take $\rho > \sigma + 1 > 0;\quad a > 0; b > 0;\quad \xi \geq 0, \delta \geq 0;\quad \lambda \geq 0;\quad \eta \geq 0;$ *and* $\mu_i \geq o, \upsilon_i \geq 0; (\forall i = 1,2,...,r)$ *then the following integration hold*

$$\int_0^\infty y^\sigma (a+by)^{-\rho} e^{\delta y^\xi}\, {}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda (a+by)^\eta\right]\times$$
$$H^{M,N;m_1,n_1;...;m_r,n_r}_{P,Q;p_1,q_1;...;p_r,q_r}\left[\begin{matrix} c_1 z_1 y^{\mu_1}(a+by)^{-\upsilon_1} \\ \vdots \\ c_r z_r y^{\mu_r}(a+by)^{-\upsilon_r}\end{matrix}\right] dx = \frac{a^{\sigma-\rho+1}}{b^{\sigma+1}} e^{\delta\left(\frac{a}{b}\right)^\xi}\, {}_2F_1\left(\alpha,\beta;\gamma;\frac{ca^{\lambda+\eta}}{b^\lambda}\right)$$
$$\sum_{k=o}^\infty\sum_{u=0}^\infty H^{M,N+2;m_1,n_1;...;m_r,n_r}_{P+2,Q+1;p_1,q_1;...;p_r,q_r}\left[\begin{matrix} a^{\mu_1-\upsilon_1} b^{-\mu_1} c_1 z_1 \\ a^{\mu_r-\upsilon_r} b^{-\mu_r} c_r z_r\end{matrix}\middle|\begin{matrix}\begin{pmatrix}2+\sigma+k\xi+u\lambda+u\eta-\rho; \\ \upsilon_1-\mu_1,...,\upsilon_r-\mu_r\end{pmatrix}; \\ \left(b_j,B_j^{(1)},B_j^{(2)},...,B_j^{(r)}\right)_{1,Q};\end{matrix}\right. \tag{13}$$
$$\left.\begin{matrix}\begin{pmatrix}-\sigma-\lambda u-k\xi; \\ \mu_1,...,\mu_r\end{pmatrix};\left(a_j;A_j^{(1)},A_j^{(2)},...A_j^{(r)}\right)_{2,P+1};\left(c_j^{(1)},C_j^{(1)}\right)_{1,p_1};...;\left(c_j^{(r)},C_j^{(r)}\right)_{1,p_r} \\ (1+\sigma+k\xi+u\lambda;\mu_1,...,\mu_r);\left(d_j^{(1)},D_j^{(1)}\right)_{1,q_1};...;\left(d_j^{(r)},D_j^{(r)}\right)_{1,q_r}\end{matrix}\right]$$

The above integral will be convergence for condition (6), (7) and (8). The integrals (13) to (15) can be proved on lines similar to those of integral (12)

3. PARTICULAR CASES:

If we take $r = 2, \eta = 0$ and $b = 1$ in eq. (10), (11), (12) and (13) then Eq. (10) reduce to (14) as follows

$$\int_0^\infty y^\sigma (a+y)^{-\rho} e^{\delta y^\xi}\, {}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda\right] H^{M,N;m_1,n_1;m_2,n_2}_{P,Q;p_1,q_1;p_2,q_2}\left[\begin{matrix} c_1 z_1 y^{\mu_1}(a+y)^{\upsilon_1} \\ c_2 z_2 y^{\mu_2}(a+y)^{\upsilon_2}\end{matrix}\right] dx$$
$$= a^{\sigma-\rho+1} e^{\delta a^\xi}\, {}_2F_1\left(\alpha,\beta;\gamma;ca^\lambda\right)\sum_{k=o}^\infty\sum_{u=0}^\infty H^{M+1,N+1;m_1,n_1;m_2,n_2}_{P+2,Q+1;p_1,q_1;p_2,q_2}\left[\begin{matrix} a^{\mu_1+\upsilon_1} c_1 z_1 \\ a^{\mu_2+\upsilon_2} c_r z_r\end{matrix}\middle|\right. \tag{14}$$
$$\left.\begin{matrix}\begin{pmatrix}-\sigma-u\lambda-k\xi; \\ \mu_1,\mu_2\end{pmatrix};\left(a_j;A_j^{(1)},A_j^{(2)}\right)_{2,P+1};\left(\rho;\upsilon_1,\upsilon_2\right);\left(c_j^{(1)},C_j^{(1)}\right)_{1,p_1};\left(c_j^{(2)},C_j^{(2)}\right)_{1,p_2} \\ \begin{pmatrix}\rho-\sigma-u\lambda-k\xi-1; \\ \upsilon_1+\mu_1,\upsilon_2+\mu_2\end{pmatrix};\left(b_j,B_j^{(1)},B_j^{(2)}\right)_{1,Q+1};\left(d_j^{(1)},D_j^{(1)}\right)_{1,q_1};\left(d_j^{(2)},D_j^{(2)}\right)_{1,q_r}\end{matrix}\right]$$

Eq. (11) reduce to (15) as follows

$$\int_0^\infty y^\sigma (a+y)^{-\rho} e^{\delta y^\xi}\, {}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda\right] H^{M,N;m_1,n_1;m_2,n_2}_{P,Q;p_1,q_1;p_2,q_2}\left[\begin{matrix} c_1 z_1 y^{-\mu_1}(a+y)^{-\upsilon_1} \\ c_2 z_2 y^{-\mu_2}(a+y)^{-\upsilon_2}\end{matrix}\right] dx$$
$$= a^{\sigma-\rho+1} e^{\delta a^\xi}\, {}_2F_1\left(\alpha,\beta;\gamma;ca^\lambda\right)\sum_{k=o}^\infty\sum_{u=0}^\infty H^{M+1,N+1;m_1,n_1;m_2,n_2}_{P+2,Q+1;p_1,q_1;p_2,q_2}\left[\begin{matrix} a^{-\mu_1-\upsilon_1} c_1 z_1 \\ a^{\mu_2+\upsilon_2} c_2 z_2\end{matrix}\middle|\right.$$
$$\left.\begin{matrix}\begin{pmatrix}2+\sigma+k\xi+u\lambda-\rho; \\ \mu_1+\upsilon_1,\mu_2+\upsilon_2\end{pmatrix};\left(a_j;A_j^{(1)},A_j^{(2)}\right)_{2,P+1};\left(c_j^{(1)},C_j^{(1)}\right)_{1,p_1};\left(c_j^{(2)},C_j^{(2)}\right)_{1,p_2} \\ \begin{pmatrix}1+\sigma+k\xi+u\lambda; \\ \mu_1,\mu_2\end{pmatrix};\left(b_j,B_j^{(1)},B_j^{(2)}\right)_{1,Q+1};\begin{pmatrix}1-\rho; \\ \upsilon_1,..,\upsilon_r\end{pmatrix};\left(d_j^{(1)},D_j^{(1)}\right)_{1,q_1};\left(d_j^{(2)},D_j^{(2)}\right)_{1,q_r}\end{matrix}\right] \tag{15}$$

Eq. (12) reduce to (16) as follows

$$\int_0^\infty y^\sigma (a+y)^{-\rho} e^{\delta y^\xi}\, {}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda\right] H^{M,N;m_1,n_1;m_2,n_2}_{P,Q;p_1,q_1;p_2,q_2}\left[\begin{matrix} c_1 z_1 y^{-\mu_1}(a+y)^{\upsilon_1} \\ c_2 z_2 y^{-\mu_2}(a+y)^{\upsilon_2}\end{matrix}\right] dx$$
$$= a^{\sigma-\rho+1} e^{\delta a^\xi}\, {}_2F_1\left(\alpha,\beta;\gamma;ca^\lambda\right)\sum_{k=o}^\infty\sum_{u=0}^\infty H^{M+1,N+1;m_1,n_1;m_2,n_2}_{P+2,Q+1;p_1,q_1;p_2,q_2}$$
$$\left[\begin{matrix} a^{\upsilon_1-\mu_1} c_1 z_1 \\ a^{\upsilon_2-\mu_2} c_2 z_2\end{matrix}\middle|\begin{matrix}(2+\sigma+k\xi+u\lambda-\rho;\mu_1-\upsilon_1,\mu_2-\upsilon_2); \\ (1+\sigma+k\xi+u\lambda;\mu_1,\mu_2);\end{matrix}\right. \tag{16}$$
$$\left.\begin{matrix}\left(a_j;A_j^{(1)},A_j^{(2)}\right)_{2,P+1};\left(\rho;\upsilon_1,..,\upsilon_r\right);\left(c_j^{(1)},C_j^{(1)}\right)_{1,p_1};\left(c_j^{(2)},C_j^{(2)}\right)_{1,p_2} \\ \left(b_j,B_j^{(1)},B_j^{(2)}\right)_{2,Q+1};\left(d_j^{(1)},D_j^{(1)}\right)_{1,q_1};\left(d_j^{(2)},D_j^{(2)}\right)_{1,q_2}\end{matrix}\right]$$

Eq. (13) reduce to (17) as follows

$$\int_0^\infty y^\sigma (a+y)^{-\rho} e^{\delta y^\xi}\, {}_2F_1\left[\alpha,\beta;\gamma;cy^\lambda\right] H^{M,N;m_1,n_1;m_2,n_2}_{P,Q;p_1,q_1;p_2,q_2}\left[\begin{matrix} c_1 z_1 y^{\mu_1}(a+y)^{-\upsilon_1} \\ c_2 z_2 y^{\mu_2}(a+y)^{-\upsilon_2}\end{matrix}\right] dx$$
$$= a^{\sigma-\rho+1} e^{\delta a^\xi}\, {}_2F_1\left(\alpha,\beta;\gamma;ca^\lambda\right)\sum_{k=o}^\infty\sum_{u=0}^\infty H^{M,N+2;m_1,n_1;m_2,n_2}_{P+2,Q+1;p_1,q_1;p_2,q_2}$$
$$\left[\begin{matrix} a^{\mu_1-\upsilon_1} c_1 z_1 \\ a^{\mu_2-\upsilon_2} c_2 z_2\end{matrix}\middle|\begin{matrix}(2+\sigma+k\xi+u\lambda-\rho;\upsilon_1-\mu_1,\upsilon_2-\mu_2);(-\sigma-\lambda u-k\xi;\mu_1,\mu_2); \\ \left(b_j,B_j^{(1)},B_j^{(2)}\right)_{1,Q};(1+\sigma+k\xi+u\lambda;\mu_1,\mu_2);\end{matrix}\right. \tag{17}$$
$$\left.\begin{matrix}\left(a_j;A_j^{(1)},A_j^{(2)}\right)_{2,P+1};\left(c_j^{(1)},C_j^{(1)}\right)_{1,p_1};\left(c_j^{(2)},C_j^{(2)}\right)_{1,p_2} \\ \left(d_j^{(1)},D_j^{(1)}\right)_{1,q_1};\left(d_j^{(2)},D_j^{(2)}\right)_{1,q_2}\end{matrix}\right]$$

4. CONCLUSION:

The H-function of several complex variables, is quite basic in nature. Therefore, on specializing the parameters of this function, we may obtain various other special functions such as Meijer's G-function, Fox's H-function, Wright's generalized hypergeometric function, Wright's generalized Bessel function, Whittaker function, generalized hypergeometric function, Mac-Robert's E-function, modified Bessel function, Bessel function of first kind, binomial function, exponential function, etc. as its special cases, and therefore, various unified integrals can be obtained as special cases of our results.

This template, modified in MS Word 2007 and saved as a "Word 97-2003 Document" for the PC, provides authors with most of the formatting specifications needed for preparing electronic versions of their papers. All standard paper components have been specified for three reasons: (1) ease of use when formatting individual papers, (2) automatic compliance to electronic requirements that facilitate the concurrent or later production of electronic products, and (3) conformity of style throughout a conference proceeding. Margins, column widths, line spacing, and type styles are built-in; examples of the type styles are provided throughout this document and are identified in italic type, within parentheses, following the example.

(This work is licensed under a Creative Commons Attribution 4.0 International License.)

International Journal of Engineering Research & Technology (IJERT)
ISSN: 2278-0181
Volume 13, Issue 12, December 2024

Some components, such as multi-leveled equations, graphics, and tables are not prescribed, although the various table text styles are provided. The formatter will need to create these components, incorporating the applicable criteria that follow.

REFERENCES

[1] Ashiq Hussain Khan, Neelam Pandey, Integrals Involving H-function of Several Complex Variables, International Journal of Scientific and Research Publications, Volume 7, Issue 2, February 2017.
[2] Bateman Project, (1953) Higher Transcendental Function, Vol. I.
[3] Gradshteyn, I.S. and Ryzhik, I.M., (2007), "Table of Integrals, Series and Products", Edited by A. Jeffrey, Daniel Zwillinger, Academic Press in an imprint of Elsevier,
[4] Mathai, A. M. and Saxena, R. K. "The H-function with Application in Statistics and Other Disciplines", Wiley Eastern Limited, New Delhi, Bangalore, Bombay, (1978).
[5] Srivastava, H. M. and Panda, R., "Some expansion theorems and generating relations for the H -function of several complex variables. I and II, Comment. Math. Univ. St. Paul. 24, fasc. 2, 119-137 (1975); ibid. 25, fasc. 2, 167-197.
[6] Panda, R. & Srivastava, H. (1976). Some bilateral generating functions for a class of generalized hypergeometric polynomials. Journal für die reine und angewandte Mathematik, 976 (283-284), 265-274. https://doi.org/10.1515/crll.1976.283-284.265
[7] Rainville, E. D., (1971), "Special Functions", Chelsea Publication Company, Bronx, New York.
[8] Srivastava, H. M., K.C. Gupta and S. P. Goyal, (1982), "The H-Functions of One and Two Variables with Applications", South Asian Publishers, New Delhi, Madras. https://doi.org/10.1002/asna.2113060427.
[9] Srivastava, H.M., (1972), "A contour integral involving fox's H-function", Indian J. Math.

(This work is licensed under a Creative Commons Attribution 4.0 International License.)